

# Baire-Class 1 and Continuity

Joel H. Shapiro

June 29, 2021

Notes for a 4/26/19 talk in the Portland State University Analysis Seminar on the continuity of derivatives, and more generally functions of Baire-Class 1, that followed up on Pieter VandenBerge's 4/12/19 talk [VB] on the possible discontinuities of derivatives. Pieter showed that every derivative has the "Intermediate Value Property;" Here we'll see that every derivative, and more generally, every function of Baire Class 1, is continuous on a dense  $G_\delta$  set.

## 1 Derivatives and Baire-Class 1

OUR SETTING will be a closed interval  $I$  of the real line  $\mathbb{R}$ . This interval could, for example, be the whole line, some half-line, e.g.  $[0, \infty)$ , or a finite interval, e.g.,  $[0, 1]$ .<sup>1</sup>

A DERIVATIVE on  $I$  is a function  $f: I \rightarrow \mathbb{R}$  for which there exists a differentiable  $F: I \rightarrow \mathbb{R}$  with  $F' = f$  on  $I$ .

THE INTERMEDIATE-VALUE PROPERTY (IVP) We say a function  $I \rightarrow \mathbb{R}$  "has the IVP" if any real number between two values of  $f$  is also a value of  $f$ .

DARBOUX'S THEOREM.[Da (1875)] *Every derivative on  $I$  has the IVP.*

We learn early on in Calculus that every *continuous* function  $I \rightarrow \mathbb{R}$  has the IVP. Might this signal that derivatives possess some vestige of continuity? We'll address this question next.

BAIRE-CLASS 1 is the collection of functions that are *pointwise limits of continuous functions*, i.e., all functions  $f: I \rightarrow \mathbb{R}$  for which there exists a sequence  $(f_n)$  of functions continuous on  $I$  with

$$f(x) = \lim_{n \rightarrow \infty} f_n(x) \quad \forall x \in I.$$

**Proposition 1.** *Every derivative on  $I$  is of Baire-Class 1.*

*Proof.* For  $x \in I$ :

$$f(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{n \rightarrow \infty} n \underbrace{\left[ f\left(x + \frac{1}{n}\right) - f(x) \right]}_{f_n(x)}$$

where each  $f_n$  is continuous on  $I$ . □

In these notes we'll prove

BAIRE'S CONTINUITY THEOREM<sup>2</sup> *Every function of Baire-Class 1 on  $I$  (in particular: every derivative) is continuous on a dense subset of  $I$ .*

<sup>1</sup> Except for our discussion of derivatives,  $I$  could even be a complete metric space.

The IVP need not have anything to do with continuity; there are functions having the IVP that take every real value in every interval. A particularly interesting example is "Conway's base 13 function," see e.g., [Om] for this, and for an interesting discussion of the IVP

<sup>2</sup> [Ba (1899)]. This is not standard terminology.

**Corollary 2.** *Dirichlet's function ( $\equiv 1$  on the rationals and 0 on the irrationals) is not of Baire-Class 1 on any interval.*

*Proof.* Dirichlet's function has no points of continuity. □

## 2 Uniform Approximation

UNLESS OTHERWISE NOTED,  $f: I \rightarrow \mathbb{R}$  is of Baire-class 1, i.e., there exists a sequence  $(f_n)$  of functions continuous on  $I$  such that

$$f(x) = \lim_n f_n(x) \quad \forall x \in I. \quad (1)$$

If the above convergence were *uniform* on  $I$ , then  $f$  would inherit the continuity of the  $f_n$ 's at *every point* of  $I$ .

Now uniform convergence of  $f_n$  to  $f$  on  $I$  means that for each  $\varepsilon > 0$  the continuous function  $f_n$  provides, for each  $n$  sufficiently large, an  $\varepsilon$ -approximation to  $f$  that is *uniform over*  $I$ . Does mere *pointwise* convergence bestow upon  $(f_n)$  some kind of uniform approximation?

To study this question, let's define, for *any* sequence of functions  $(f_n): I \rightarrow \mathbb{R}$ , and any  $\varepsilon > 0$ , its set of " $\varepsilon$ -Cauchy-ness"<sup>3</sup>

$$C(\varepsilon) = \bigcup_{N>0} \underbrace{\bigcap_{n,m \geq N} \{|f_n - f_m| \leq \varepsilon\}}_{=: F_N(\varepsilon)}. \quad (2)$$

<sup>3</sup> Once again: not standard terminology.

However the abbreviation  $\{g \leq \varepsilon\}$  is standard, and stands for the set  $\{x \in I: g(x) \leq \varepsilon\}$ , with similar notation for other inequalities and equalities involving functions.

Should each  $f_n$  be *continuous* on  $I$ , the sets in curly brackets on the right-hand side of equation (2) will all be closed in  $I$ , hence so will each set  $F_N(\varepsilon)$ . Thus each  $C(\varepsilon)$ , while not necessarily itself closed, will be a *countable union* of closed sets (an " $F_\sigma$ -set" in the now-standard terminology introduced in 1899 by Baire [Ba]).

RETURNING NOW to the situation described by equation (1), we have, from the fact that  $(f_n(x))$  is, for each  $x \in I$ , a Cauchy sequence of real numbers:

$$I = C(\varepsilon) = \bigcup_{N \in \mathbb{N}} F_N(\varepsilon) \quad \forall \varepsilon > 0. \quad (3)$$

Since each  $F_n(\varepsilon)$  is a closed subset of  $I$ , this sets the stage for:

**The Baire Category Theorem** ([Ba],1899). *If  $I$  is a countable union of closed subsets, then at least one of these must contain an interval.*

We'll defer the proof of this famous result to §5, after we've come to appreciate it's role in unravelling the secrets of Baire-Class 1.

**Corollary 3.** *Suppose  $f$  is a function of Baire-class 1, as described by equation (1) above. Then for every  $\varepsilon > 0$  there exists a positive integer  $N = N(\varepsilon)$  and an open interval  $J = J(\varepsilon)$  such that for every  $x \in J$ :*

- (a)  $|f_n(x) - f_m(x)| \leq \varepsilon$  for each  $n, m \geq N$ , and  
 (b)  $|f_N(x) - f(x)| \leq \varepsilon$  for every  $n \geq N$ .

*Proof.* Part (a) follows directly from the Baire Category Theorem and the representation (3) of  $I$  as a countable union of closed sets. Part (b) follows from part (a) upon setting  $m = N$  and letting  $n \rightarrow \infty$ .  $\square$

**SUMMARY** for a Baire-Class 1 function as described by equation (1):

*For every  $\varepsilon > 0$  there exists  $N = N(\varepsilon) \in \mathbb{N}$  such that  $|f - f_N| < \varepsilon$  on some open interval  $J(\varepsilon)$  of  $I$ .*

Our next task will be to understand what degree of continuity (if any) the “ $\varepsilon$ -uniform approximator”  $f_N$  passes on to its “approximatee”  $f$ . Here “degree of continuity” will be measured by the notion of “oscillation.”

### 3 Oscillation

Suppose  $S$  is any set and  $f: S \rightarrow \mathbb{R}$ . The *oscillation* of  $f$  over  $S$  is defined as

$$\Omega_f(S) = \sup_{x, y \in S} |f(x) - f(y)|. \quad (4)$$

Note that  $\Omega_f(S)$  decreases with  $S$ , i.e.,

$$S_1 \supset S_2 \implies \Omega_f(S_1) \geq \Omega_f(S_2).$$

It follows that: for an open interval  $J$  a function  $f: J \rightarrow \mathbb{R}$ , and a point  $x \in J$ , the limit

$$\omega_f(x) = \lim_{r \rightarrow 0^+} \Omega_f(J(x, r)) \quad (5)$$

Here  $J(x, r)$  is the open interval  $(x - r, x + r)$ .

exists and is  $\geq 0$ . We call  $\omega_f(x)$  the *oscillation of  $f$  at  $x$* . Here are some of its crucial properties; the proofs of which are all easy exercises.

**Proposition 4.** *For a real-valued function  $f$  on an open interval  $J$ :*

- (a) *If  $|f| \leq M$  on  $J$  then  $\omega_f \leq 2M$  on  $J$ .*  
 (b)  *$f$  is continuous at a point  $x \in J$  if and only if  $\omega_f(x) = 0$ .*  
 (c) *For each  $\lambda > 0$ , the set  $\{\omega_f < \lambda\}$  is open in  $J$ .*  
 (d) *If both  $f$  and  $g$  are real-valued functions on  $J$ , then at each point of  $J$*

$$\omega_{f+g} \leq \omega_f + \omega_g.$$

**Corollary 5.** *Suppose:*

- $J$  is an open real interval,
- $g: J \rightarrow \mathbb{R}$  is a function that is continuous on  $J$ ,
- $\varepsilon$  is a positive real number, and
- $f$  is any real-valued function on  $J$  with  $|f(x) - g(x)| \leq \varepsilon$  for each  $x \in J$ .

Then:  $\omega_f(x) \leq 2\varepsilon$  for every  $x \in J$ .

*Proof.* Fix  $x \in J$ . Then from Proposition 4(d):

$$\omega_f(x) = \omega_{(f-g)+g}(x) \leq \omega_{f-g}(x) + \omega_g(x).$$

Now  $\omega_g(x) = 0$  from the continuity of  $g$  and Proposition 4(b), while by part (a) of that Proposition and the uniform estimate on  $|f - g|$  we have  $\omega_{f-g}(x) \leq 2\varepsilon$ .  $\square$

#### 4 Proof of Baire's Continuity Theorem

RECALL THAT WE'RE GIVEN a closed interval  $I$  of the real line and a real-valued function  $f$  on  $I$  that's the pointwise limit of a sequence  $(f_n)$  of functions that are continuous on  $I$ .

TO SHOW:  $f$  is continuous on a dense subset of  $I$ .

To this end, fix  $\varepsilon > 0$  and let  $G_\varepsilon = \{x \in I: \omega_f(x) < \varepsilon\}$ . By part (c) of Proposition 4  $G_\varepsilon$  is open in  $I$ .

CLAIM:  $G_\varepsilon$  is dense in  $I$ .

Indeed: we know from Corollary 3 that there is an interval  $J \subset I$  and an index  $N \in \mathbb{N}$  such that  $|f - f_N| \leq \varepsilon/3$  on  $J$ . Thus  $\omega_f$  is  $\leq 2\varepsilon$  at every point of  $J$ , hence  $J \subset G_\varepsilon$ . In particular, this shows (finally!) that  $G_\varepsilon$  is non-empty.

HOWEVER THE CLAIM demands that we show  $G_\varepsilon$  has non-empty intersection with *every* open subinterval  $H$  of  $I$ , not just  $J$  (which it contains). No problem: apply the result of the last paragraph, with  $\overline{H}$  (the closure of  $H$ ) in place of  $I$ , and  $G_\varepsilon \cap H$  (which is open and dense in  $\overline{H}$ ) in place of  $G_\varepsilon$ . The result is that  $G_\varepsilon \cap H$  is not empty, as desired.

Now the set of points of continuity of  $f$  (indeed, for any function on  $I$ ) is

$$\{\omega_f = 0\} = \bigcap_{n \in \mathbb{N}} \{\omega_f < 1/n\} = \bigcap_n G_{1/n}, \quad (6)$$

a countable intersection of dense, open subsets of  $I$  which, by an "alternate version" of the Baire Category Theorem (see next section) is dense.  $\square$

IN ADDITION TO his designation “ $F_\sigma$ ” for countable unions of closed sets, Baire introduces the term “ $G_\delta$ ” for countable *intersections of open sets*. Thus Eqn. (6) shows that for any real-valued function  $f$  on  $I$ :

*The set of points of continuity is a  $G_\delta$  set.*

... which, in this generality, may be empty.

## 5 Proof of Baire’s Category Theorem

WE’VE EMPLOYED two equivalent versions of the Baire Category Theorem (henceforth, the “BCT”), which we’ll attack in its natural setting: that of metric spaces  $(X, d)$ .<sup>4</sup>

<sup>4</sup> Feel free to restrict attention to the metric space most comfortable for you, e.g.  $X = \mathbb{R}$  or some closed interval therein, and  $d =$  the usual length function on  $\mathbb{R}$

**The Original BCT.** *If a complete metric space is the countable union of closed subsets, then at least one of these sets has nonvoid interior.*

**The Alternate BCT.** *In a complete metric, every countable intersection of dense open sets is dense.*

WE’LL PROVE the equivalence of these two versions, after which we’ll establish the “alternate” one.

**Proof that “Original BCT” implies “Alternate BCT.”** Assuming the Original BCT, suppose  $\{G_n\}$  is a countable family of dense, open subsets of our metric space  $X$ .

TO SHOW:  $G = \bigcap_n G_n$  is dense, i.e.,  $G$  has nontrivial intersection with every nonvoid open subset of  $X$ .

Since each  $G_n$  is open, its complement  $F_n = X \setminus G_n$  is closed, and since  $G_n$  is dense,  $F_n$  is nowhere dense, i.e., dense in no open set.<sup>5</sup> By the Original BCT,  $X \neq \bigcup_n F_n = X \setminus \bigcap_n G_n$ . Thus  $G = \bigcap_n G_n$  is not empty.

<sup>5</sup> If  $F_n$  were dense in some open subset  $V$  of  $X$ , then since  $F_n$  is closed,  $V$  would be contained in  $F_n$ . But then no point of  $V$  could be a limit point of  $G_n$ , hence  $G_n$  could not be dense.

We’re now in a situation similar to the one encountered in our proof of Baire’s Continuity Theorem. We want to show that  $G$  intersects every nonvoid open set, but have just shown that it’s nonempty! Once again, generality saves the day!

Let  $V$  be a nonvoid open subset of  $X$ , and replace  $X$  by  $\bar{V}$ , the closure of  $V$  in  $X$ . Then  $\bar{V}$  is a complete metric space (in the metric of  $X$ ), in which each  $G_n \cap V$  is a dense open subset. It follows from the last paragraph that  $G \cap V = \bigcap_n (G_n \cap V)$  is nonempty, i.e., that  $G$  intersects  $V$ . □

**Proof that “Alternate BCT” implies “Original BCT.”** Suppose now that we’ve proved the alternate BCT. Suppose  $\{F_n\}$  is a countable collection of closed subsets of  $X$  with  $X = \bigcup_n F_n$ . Then  $G_n = X \setminus F_n$  is open in  $X$  for each  $n \in \mathbb{N}$ , and  $\bigcap_n G_n$  is empty. By the alternate form of the BCT, some  $G_n$  must fail to be dense, hence the corresponding complement  $F_n$  must have nonvoid interior (i.e., it must contain a nonvoid open set). □

BEFORE PROVING THE ALTERNATE BCT, let's take a moment to appreciate the special nature of dense open sets. We know that two dense sets may be disjoint (e.g., the rationals and the irrationals in the real line), and the same can happen for two open sets (e.g. the intervals  $(0,1)$  and  $(1,2)$  of  $\mathbb{R}$ ). But this can't happen for two sets that are *both dense and open*.

Indeed, in a metric space, suppose  $V_1$  and  $V_2$  are both dense and open, and let  $U$  be any nonvoid open subset of our space. Since  $V_1$  is dense, it has a nonvoid intersection with  $U$ . Since  $V_1 \cap U$  is open and non-void, and  $V_2$  is dense, it has a non-void intersection with  $V_1 \cap U$ . Thus  $V_1 \cap V_2$  has non-void intersection with any nonvoid open set  $U$ , and is therefore dense in our space.

More generally, an easy induction shows that:

*In any metric space, the intersection of a finite family of dense open sets is dense.*

THE POINT OF the "Alternate BCT" is that for *complete* metric spaces this result extends to *countable* families of dense open sets. The argument will be similar to the one you just used to show that intersections of finite collections of dense open sets are dense—but now it requires a little more precision

EXERCISE. In an *incomplete* metric space this result can fail.

**Proof of the Alternate BCT.** Our setting is a metric space  $(X, d)$  that is *complete*, i.e., in which every Cauchy sequence converges.

Let's fix a countable collection  $\{V_n\}$  of subsets of  $X$ , each of which is open and dense in  $X$ , and set  $G = \bigcap_n V_n$ .

TO SHOW:  $G$  is dense in  $X$ .

Since  $V_1$  is open and nonvoid it contains a ball  $B_1$  whose closure  $\overline{B_1}$  also lies in  $V_1$ . Since  $V_2$  is dense and open, its intersection with  $B_1$  is open and nonvoid, hence  $V_2 \cap B_1$  contains a ball  $B_2$  whose closure also lies in  $V_2 \cap B_1$ , and has diameter  $\leq 1/2 \times$  the diameter of  $\overline{B_1}$ . Continuing by induction we produce a decreasing sequence  $\{\overline{B_n}\}$  of closed balls, with  $\overline{B_n} \subset V_n$  and the diameter of  $\overline{B_n} \leq 1/2^{(n-1)} \times$  that of  $\overline{B_1}$ . By Cantor's Nested Set Theorem,  $\bigcap_n \overline{B_n} \neq \emptyset$ , and by our construction, any point of this intersection must lie in every  $V_n$ . Thus  $\bigcap_n V_n \neq \emptyset$ .

Our goal is to prove that  $G$  is dense in  $X$ , but so far we just know it's non-empty. To complete the proof, fix a nonempty open subset  $U$  of  $X$  and apply the result just obtained, with  $X$  replaced by the closure  $\overline{U}$  of  $U$  and  $V_n$  replaced by  $V_n \cap U$ . The result—thanks to the fact that every closed subset of a complete metric space is itself complete (in the original metric)—is that  $G \cap U = \bigcap_n (V_n \cap U)$  is nonempty, which establishes the density of  $G$  in  $X$ .  $\square$

**CANTOR'S INTERSECTION THEOREM.** *In a complete metric space, every countable decreasing sequence of non-empty closed sets with diameter  $\rightarrow 0$  has non-empty intersection.*

*Proof.* Suppose  $(F_n)$  is our decreasing sequence of closed sets. By passing to an appropriate subsequence, we may assume that the diameter of  $F_n$  is  $\leq 1/2^n$ . Fix  $x_n \in F_n$ . Then for  $m < n$  the triangle inequality yields

$$d(x_m, x_n) \leq \sum_{j=m}^{n-1} d(x_j, x_{j+1}) \leq \sum_{j=m}^{n-1} \frac{1}{2^j}.$$

CONCLUSION:  $(x_n)$  is Cauchy in  $X$ , hence convergent.<sup>6</sup> □

<sup>6</sup> The proof shows that  $\bigcap_n F_n$  is a singleton.

## 6 Concluding remarks

### 6.1 Baire spaces.

A *Baire space* is a topological space for which the intersection of every countable family of dense, open sets is dense. Thus: *Every complete metric space is a Baire space.*

Although completeness is not preserved by homeomorphisms, “Baire-ness” is preserved. Thus:

*Every topological space homeomorphic to a complete metric space is a Baire space.*

EXAMPLE: the open unit interval, which is not complete in the metric of the real line, is nonetheless homeomorphic to the line

In particular:

*The conclusion of the Baire Category Theorem holds for every open interval of the real line.*

Note also that our proof of the Baire Category Theorem depended only upon the fact that in a complete metric space, every decreasing sequence of nonempty closed sets has nonempty intersection. Thus:

*Every compact topological space is a Baire space.*

### 6.2 Dense $G_\delta$ sets are (usually) uncountable!

For definiteness, let's take as our initial setting the real line. Suppose  $G$  is a dense  $G_\delta$  subset of  $\mathbb{R}$ . Thus  $G = \bigcap_n V_n$  where each  $V_n$  is an open subset that is itself dense in  $\mathbb{R}$ . To see that  $G$  is uncountable, suppose—for the sake of contradiction—that it is not, i.e., suppose that  $G = \{x_1, x_2, \dots\}$ . Then  $W_n = V_n \setminus \{x_1, x_2, \dots, x_n\}$  is open and dense in  $\mathbb{R}$ , and

$$\bigcap_n W_n = G \setminus \{x_n\}_1^\infty = G \setminus G = \emptyset,$$

contrary to the Baire Category Theorem. □

THE ARGUMENT ABOVE works, word-for-word, to establish:

**Proposition.** *In any Baire space without isolated points, every dense  $G_\delta$  set is uncountable.*

### 6.3 Back to the beginning.

Let's return to our original closed interval  $I$  of the real line. We've shown that for every real-valued function on  $I$ , its set of points of continuity is a (possibly empty)  $G_\delta$ . However, if our function is of Baire-Class 1 we've shown that its  $G_\delta$ -set of continuity is *dense*. In view of what we've just seen in §6.2, we now know that this set is, in addition, *uncountable*.

### References

**Ba** Baire, Rene-Louis, *Sur les fonctions de variables réelles*, Ann di Mat. 3 (1899) 1-123.

**Da** Darboux, Jean Gaston, *Mémoire sur les fonctions discontinues*, Ann. Sci. E.N.S. (2) 4 (1875) 57-112.

**Ox** Oxtoby, John C. *Measure and Category*, Springer 1971.

**Om** Oman, Greg *The converse of the Intermediate Value Theorem . . .*, Missouri J. Math. Sci. 26 (2014) 134-150.

**Ru** Rudin, Walter. *Real and Complex Analysis*, 3rd edition, McGraw-Hill 1987.

**VB** VandenBerge, Pieter. *Derivatives and their discontinuities*, freely available at: [https://joelshapiro.org/Pubvit/Downloads/vandenberge\\_derivatives.pdf](https://joelshapiro.org/Pubvit/Downloads/vandenberge_derivatives.pdf)

Here you'll find not only Darboux's IVP theorem for derivatives, but also as his introduction of the "Darboux integral" (based on the limit of upper and lower integrals).